

ARGEMONE MEXICANA L.: A WEED HAVING NUMEROUS THERAPEUTIC AND PHARMACEUTICAL USES

Shinde, Y. P.,¹ and Pachkore, G. L.²

Received: 31/3/2023

Revised: 25/5/2023

Accepted:26/5/2023

Published: 30/12/2023

Abstract

Many plants in the enormous floral diversity are considered weeds or wild and have no commercial value. Hence, some what ignored. One such weed that is wildly proliferating over Kalsubai Region of western ghat is *Argemone mexicana* L. Due to its antagonist, this herb may survive without any special care. Genetic capacity to synthesise protective phytochemicals under pressure. These phytochemicals also serve a variety of other purposes, including medicine. This review has highlighted numerous notable medical applications of this herb as well as some of its negative consequences.

Keywords: Preserve, Adaptations, Blossoming

Introduction

There are an infinite number of plants on this green planet that are regarded as undesired or weeds since they have no apparent economic value. Some plants do, however, thrive in harsh environments and have made good adaptations to them. But, nothing that Mother Nature has made is worthless; everything has value. Hence, these herbaceous weeds also have something useful in their genomic makeup, but to be properly investigated in a sustainable way, they need keen eyes and considerable passion. Several plants that were once thought to be useless have now been shown to be extremely useful in terms of their phytochemicals. As the wild shrubs, trees, and herbs are truly major sources of new phytochemicals, one goal of biodiversity study is to preserve these priceless gene pools.

Argemone mexicana (AM), a perennial weed that is endemic to Mexico and primarily associated with severe settings and agricultural crops, is a member of the Papaveracea family. That is a top weed for lots of money. of crops in the hu-

mid temperate, subtropical, and tropical regions of the world. This plant is known by a variety of common names, including Mexico prickly poppy, blossoming thistle, Cardo/Cardosanto, etc. This plant is known by a number of names in India, including Agara, Bharband, Bharbhar, Brahamadandi, Kantakusama, and PeelaKanteela.

Because it produces severe itching and annoys farmers, this fussy plant species is seen as a risk to health if consumed with tainted food. *A. mexicana* has been determined to be dangerous because, if unintentionally swallowed, its seeds pose a threat to human or animal health (NDA, 2001).

Taxonomic rank of the plant:

Phylum: Spermatophyta;
Class: Dicotyledonae;
Order: Papaverales;
Family: Papaveraceae;
Genus: Argemone;
Species: *Argemone mexicana* L.

Historical background

¹R.G.Bagdia Arts, S.B. Lakhotia Commerce and R. Bezonji Science College, Jalna.

²VasantdadaPatil Arts Commerce & Science College, Patoda, Beed, Maharashtra India
Corresponding Author-Shinde Y P email-ypshinde129@gmail.com

A. mexicana has been inadvertently introduced through seeds as a showy plant. Apart from this some of the tribes used to cultivate this plant basically for ethnic uses (Healy, 2016; APPEAR, 2016; PROSEA, 2016), the plant is still comparatively infrequent and assumed that it may become a difficult challenge for arable land in the future. Being an annual herb with a soaring possibility of the auxiliary spread since the plant is still considered as an attractive showy plant across the world. The occurrence of species is very frequent at the sites of construction and it is also preferred for landscaping of any new landscape (Foxcroft et al., 2006). The seeds of this plant are used to disperse easily through contaminated seed products, soils and, crop (Healy, 1961; PROTA, 2016).

Taxonomic description

Argemone name was taken from the Greek word 'argena', sense 'cataract of the eye', and this name was used by the orthodox researchers, viz., Dioscorides (AD 40-90) and Pliny (AD 23-79) basically for a number of prickly poppies, the fluid of which was apparently considered as a cure for cataract. The term 'mexicana' coalesces Mexico with the Latin suffix 'ana', taken from the source nation (Parsons and Cuthbertson, 1992). *A. mexicana* is a twelvemonthly herbaceous plant, up to 120-150 cm elevation with a faintly branched tap root system. The stem is upright, pronged, generally spiny, pale to bluish-green in color (Plate 1: Figure a) and exudes an offensive smelled the yellow juice when cut. Foliages are alternate, petioles absent, roughly covering the stem, equal to 15 cm long, acutely lobed with sporadically toothed, prickly margins; veins are conspicuous, grayish-white. Flowers borne singly, 2.4- 4.6 cm in diameter, having 1-2 green bracts; calyx 3, spiny; corolla 4-6, yellow to orange, surface glabrous; stamens several (Plate 1: Figure b). Fruit is a capsule, prickly, 2.5-4.5 cm long and 2 cm broad, amid 4-6 valves with aperture at the tip to release abundant brownish-black, nearly spherical seeds having the diameter of about 1 mm, a fine network of veins present, oily in nature (Lucas, 1962).

Distribution

In India In India, this species is prevalent in states like, Andhra Pradesh, Assam, Bihar, Delhi, Gujarat, Karnataka, Madhya Pradesh, Maharashtra, Tamil Nadu, Telangana, Rajasthan, Uttar Pradesh, and West Bengal (Sharanappa and Vidyasagar, 2014).

Distribution

In India In India, this species is prevalent in states like, Andhra Pradesh, Assam, Bihar, Delhi, Gujarat, Karnataka, Madhya Pradesh, Maharashtra, Tamil Nadu, Telangana, Rajasthan, Uttar Pradesh, and West Bengal (Sharanappa and Vidyasagar, 2014).

In rest of the World This plant species is usually spread in all parts of the world. AFRICA: Nigeria, Namibia, Niger, Senegal, Seychelles, Sierra Leone, Somalia, South Africa, Sudan, Tanzania, Togo, Uganda, Zambia, Zimbabwe; ASIA: Bahrain, Bangladesh, Bhutan, Cambodia, China, Hong King, Indonesia, Java, Japan, Pakistan; MIDDLE EAST: Israel, Saudi Arabia, Turkey, Iran, Iraq, Syria; EUROPE: Italy, Spain, Switzerland, United Kingdom; NORTH AMERICA: United States of America, Anguilla, Antigua, Barbados, Belize, Canada, Hawaii; SOUTH AMERICA: Argentina, Bolivia, Brazil, Chile, Columbia, Ecuador, Guyana, Peru, Uruguay; OCEANIA: Australia, Fiji, New Zealand (Bodeker and Graz, 2013).

Habitat

A. mexicana is tailored to an extensive variety of habitats. The plant prefers the areas having a distinctive dry period (PROTA, 2016). It usually crops up as a common weed of agricultural land, pastures and in dumping yards, railways, roadsides, etc.

Affected plants by this weed

A. mexicana is a weed for the majority of cropping systems, including millets, cereals, vegetables, legumes, fibre yielding crops (sisal, cotton) and perpetual crops like coffee and sugar cane. It appears that any crop has the impending to be contaminated with *A. mexicana* if

grown within the habitat array of this plant.

Economic Impact

Apart from a principal weed of millets, cereals, cotton, vegetables, coffee, timber and fibre yielding plants, this plant is also considered as a potent contaminant in poultry and for grazing animals. The species produces aflatoxins of poisonous nature that are lethal to herbivorous animals, which even found in affected cattles' milk, eggs and their mutton based products (Alemayehu and Desalegn, 2016).

Environmental impact

Islam et al. (2003) has discovered that this weed reduces biodiversity (Kumar and Rohatgi 1999). The plant is known to produce certain allelochemicals that can affect the seed germination, subsequent growth and content of photosynthetic pigments in nearby plants in native ecosystems (Namkeleja et al. (2014).

Social impact

A. mexicana has main hammering on human wellbeing in India and adjacent nations, where the safe to consume vegetable oil either unintentionally tainted with this weed or deliberately mixed by dodgy traders. Dropsy epidemic (Delhi; 1998) is a well known example of this unlawful act (Jha et al., 2001; Sharma et al., 2002). Sharma et al. (1999) reviewed the scientific effects of adulterated oil and recommended precautionary measures. Furthermore, in the northern parts and central parts of India, this weed has been recognized as a potent allergen (Singh and Kumar, 2004).

Uses

Economic worth

A. mexicana has revealed significant antimicrobial action against the bacteria strains, viz., *Staphylococcus agalactiae*, *S. aureus* and *Escherichia coli*, with impending in the pharmaceutical production (Alemayehu and Desalegn, 2016). This weed has been cultivated for its seed oil, which is specifically used at industrial scale for soap manufacture and also for fuel production (Hanelt and IPK, 2016).

Social benefit

A. mexicana is used for spiritual reasons by many tribes (Hanelt and IPK, 2016). The extracts of the leaves, floral buds and seeds of this weed have been evaluated in laboratory conditions for their insectidal potential (Chitra et al., 1997), common crop pathogens (Singh et al., 1993) and also against nematodes (Das and Sukul, 1988; Saxena and Tabassum, 2000; Shaukat et al., 2002). Aqueous extracts were successfully assessed against, *Lipeuruslawrensis* (Kumar et al., 2002). Few researches recommended that the root extracts can be utilized to avert oviposition and work as ovicidal in opposition to *Aedesegypti* (Warikoo and Kumar, 2014). Von Weizsäckerl (1995) stated that this weed is used in India to set up an antifeedant spray similar as the foliage of *Azadirachta indica* (Neem). Plant extracts of *A. mexicana* eagerly kill the *Biomphalaria glabrata* (snail) and thus regarded as molluscicide for the fairly economical management of schistosomiasis in humans (Melendez and Capriles, 2002).

Medicinal value

Medicinal possessions have been recognized to the seed sap and its oil (Holm et al., 1977). Ethnobotanically the entire plant is used as a blend to cure asthma. The root is mixed with alcohol (rum) to cure stomach pain. The sap of the stem (cut ends) is useful for the cure for toothache. Kids having obscure urination are given mixtures of petals (DeFilippis et al., 2004). In Madhya Pradesh (India) it is designated as a homeopathic preparation (Oudhia et al., 1998). In African nations, leaves of the plant are used as a cosmetic (Rukangira, 2001). The seeds are ground and mixed with beer/tea to augment their strength (Verdcourt and Trump, 1969). In India, the minute quantity of seeds of this plant is mixed with mustard oil to increase its pungency, however, above that minute quantity the mixing of its seeds to mustard is considered as an adulteration.

Environmental services

Although this herb species has been stated as toxic to animals, however, this plant is palatable

by lemurs, which were seen consuming the stems of this weed when other resources were insufficient after a destructive cyclone in Madagascar. Those lemurs were then observed and no harmful concerns related to normal healthiness were reported by the researchers (LaFleur and Gould, 2009). Hence it can be concluded that the shoots are risk free and only seeds have toxic substances.

Ethnobotanical uses

Among many tribes this plant has certain ritual uses which are evident by their folklores (Hanelt and IPK, 2016). The different parts of this weed possess strong emetic, sedative actions and conventionally been used to take care of syphilis and various skin-diseases (Krishnamurthy, 1969; Savithamma et al., 2007). In cough and asthma seeds are given as a remedy. Seeds are also found laxative in nature with emetic, expectorant and demulcent properties. The root is an anti-helminthic (Nadkarni, 1982). Chemical characterization of this plant has discovered the existence of certain alkaloids, amino acids, phenolics and fatty acids (Hussain et al., 1983; Harborne and Williams, 1983). The plant contains several alkaloids, viz., protopine, berberine, sanguinarine, optisine, chelerytherine etc. While, the seed oil has fatty acids, viz., palmitic, myristic, oleic, linoleic acids, etc. The sap of the plant is yellow and contains minute quantities of berberine, also potassium nitrate was identified among the salts naturally existing in the plant.

Phytochemical evaluation

A huge number of phyto-constituents have been isolated and characterized from different plant parts, viz., alkaloids, phenolics, amino acids and fatty acids. Four quaternary isoquinoline alkaloids, jatrorrhizine, columbamine, dehydrocorydalmine, and oxyberberine, have been isolated from the whole plant of *A. mexicana* and their structures recognized by spectral confirmation (Papova et al., 1980; Das and Khanna 1997; Singh et al., 2010). Compounds of aliphatic nature viz., mexicanic acid and mexicanol have been isolated from leaves of *A. mexicana*. Likewise, from seeds, isoquinoline alkaloids such as

dihydropalmitine hydroxide; protopine and berberine have been isolated (Plate 1: Figure c). The seed oil contains 40% free glycerides of fatty acids (Anonymous, 2004). This plant conventionally used as an effective diuretic remedy. Furthermore, the plant exhibits anti-inflammatory, antihelminthic, injury healing, and anti-microbial properties (Bhattacharjee et al., 2006). The plant is known as an astringent, pungent, purgative, seditious, expectorant, emetic, antipyretic, ophthalmic, stomachic. The plant is remarkable in curing disease like leprosy, other skin infections, swelling, etc. (Prasanna et al., 2007). Roots are found valuable in the case of guinea-worm invasion, skin diseases, leprosy, inflammations, poisoning, digestive disorders (flatulence and constipation), colic, malaria, etc., (Sharanappa and Vidyasagar, 2014). Seeds are purgative and tranquillizer; they have a remarkable resemblance with mustard seeds and hence occasionally used to mixed with mustard seeds for unlawful monetary gain (Pahwa and Chatterjee, 1989). The seeds produce a non edible lethal oil and causes fatal dropsy disease when used with mustard oil for cooking and show lots of noxious effect. Seeds are also useful in irritable cough, asthma, pertussis, leprosy, skin diseases, injuries, dental caries, rheumatagia, constipation, flatulence and colic condition (Bhattacharjee et al., 2006). The leaves are useful in curing cough and cold, injuries, newly developed ulcers and in various skin diseases caused by microorganisms. Leaf juice is used in ophthalmology to treat ophthalmia and murkiness of the cornea (Orient Longman, 1998). The latex is useful in dropsy, leprosy, skin diseases, blisters, jaundice, conjunctivitis, burning sensation, inflammation and malaria (Kirtikar and Basu, 1991). Though the oil is known to cause dropsy yet found useful in the treatment of leprosy, skin diseases, indolent ulcers, injuries, flatulence, constipation, colic and rheumatagia. In homeopathy, a drug extracted from this plant is used to cure the trouble caused by the tape worm (Bhardwaj *et al.*, 1982). A problem known as ‘Hepatotoxicities’ is extensively came across worldwide due to the toxicity caused by this plant when used in higher doses, and considered

as one of the ten foremost causes of death globally, thus, there is a great need to perform toxicity related research to resolve the safe dose for variously stated pharmacological action. There is a need to work and identify the various isolated phyto-constituents, along with their pharmacological activity with finely calculated doses to avoid toxicity. Consequently, several researches are going on with this herb that have mainly concentrated on its hepatoprotective action, anti-microbial potential (Bhattacharjee *et al.*, 2006), anti HIV activity (Chang *et al.*, 2003a) and cytotoxic potential (Chang *et al.*, 2003b).

Major pharmacological analysis

Hepatoprotective activity

The bark of root was examined on CCl₄ (Carbon tetra Chloride) tempted liver damaged (rat model). Severe toxicity study, efficiency status, blood test and biochemical assays of the tissue (ALT, AST, total protein, glucose, bilirubin, LM and EM), *etc.*, have been evaluated for Hepato-protective act. On the basis of results it was exhibited that this herb certainly has an elevated prospective in a curative role like healing of liver parenchyma and renewal of damaged liver cells hence considered as an effective liver tonic (Pingale *et al.*, 2008).

Antimalarial activity

A decoction of *A. mexicana* was evaluated in increasing doses to find out the suitability of the decoction as healer for the treatment of malaria. It was found that the malaria patients showed enhanced ability to recover when treated with controlled doses of decoction. However, complete clearance of Plasmodium falciparum was not achieved (Willcox *et al.*, 2007).

Antibacterial activity

The basic extracts of *A. mexicana* foliage and seeds of were assessed by agar well diffusion technique to find out their potential against Staphylococcus aureus and Bacillus subtilis (Gram+ bacteria), Escherichia coli and Pseudo-

monas aeruginosa (Gram- bacteria) which are known as pathogenic and multi-drug resistant. Although, all the extracts were found efficient, hitherto the methanol extract exhibited utmost reticence against the selected bacterial strains, followed by warm aqueous extracts and cold aqueous extract, respectively (Sakthivadivel *et al.*, 2003). Likewise, the alcoholic and petroleum ether extracts of above ground parts of this herb were assessed for antimicrobial activity against, Bacillus subtilis (Gram+) and Escherichia coli (Gram-). Which showed high antibacterial activity on the bacterial strains, however the alcoholic extract showed superior action compared to the petroleum ether extract (Bhattacharjee *et al.*, 2006).

Antiplasmodial activity

A. mexicana also showed antiplasmodial activity at varying doses of extract (per kg/body wt.) About twenty species of genus Argemone have been evaluated and it was found that with the IC₅₀ values of 9- 43 mg dry extract ml⁻¹ these species possess antiplasmodial activity, *e.g.*, plant extracts [in vitro inhibition (%)] against chloroquine liable strain of *P. falciparum* (Simonsen *et al.*, 2001).

Cytotoxic activity

By fractionation of the plant's extract (in chloroform), an alkaloid was isolated and subsequently tested for its cytotoxicity to human being nasopharyngeal carcinoma and gastric cancer cell lines. Later, the alkaloid chelerythrine showed significant activity against cancerous cell line, whilst angoline was reported as potent inhibitor of cancerous cell lines (Chang *et al.*, 2003).

Larvicidal action

A. mexicana extract in petroleum ether at higher concentrations, after acetone fraction showed larvicidal properties and also growth restraining action in opposition to the subsequent (second) instar larvae of *Aedes aegypti* (Sangameswaran *et al.*, 2004).

Vasorelaxant action

The vascular effects of a methanolic extract of the above ground plant parts of *A. mexicana* was explored in aortic rings of rat. Remarkably, the extract created relaxation from contraction tempted by norepinephrine in a dose-dependent mode. At elevated doses (300 and 1000 $\mu\text{g ml}^{-1}$), the extract caused noteworthy added stress. The results showed that the extract encourages a straight dual explicit consequence upon the smooth muscle, partly mediated by adrenergic receptors (Páez-Sánchez et al., 2006).

Injury curative action

The curative effects for an injury of the foliage extract (in 50% ethanol) and latex of *A. mexicana* plants were examined in an albino rat model, using excision and incision injuries both. Topical use of the plant extract and its latex, correspondingly, conferred approximately 67 and 57% curing after 12 days in the cutting out mode and augmented tensile potency to about 190 and 155 gm in the slit out mode. Though, the plant extract and latex both were not found as successful as the customary nitrofurazone salt (Ghosh et al., 2005).

Anti-HIV action

In the methanolic extract of *A. mexicana* few alkaloids have been isolated and were evaluated for their anti-HIV action, viz., benzo[c]phenanthridine (+/-)-6-acetyldihydro chelerythrine showed noteworthy anti-HIV action in H9 lymphocytes cells with EC₅₀ and TI values of 1.77 $\mu\text{g ml}^{-1}$ and 14.6, respectively (Chang et al., 2003).

Antiasthmatic action

A. mexicana seed powder (100–200 mg) taken twice daily for 2 weeks showed noteworthy outcome on the incidence of asthma as antiasthmatic activity (Bhalke and Gosavi, 2009).

Antistress and antiallergic action

Stem of *A. mexicana* in asthma induced by leucocytosis and milk-induced eosinophilia showed antiallergic and anti-stress impending. This

showed that polar constituents of plant stem are conscientious for antistress and antiallergic activity (Piacente et al., 1997).

Molluscicidal action

Molluscicidal possessions of seeds of this weed against snail *Lymnaea acuminata* were done. It was observed that the molluscicidal action of plant's seed powder was both time and dose dependent. In seed powder, protopine and sanguinarine were recognized as the active component caused snail fatality by co-migration of a vigorous agent (Singh and Singhm 1999; Meléndez and Capriles, 2002).

Opioid withdrawal

In the methanolic plant extract, the pure compounds isolated as protopine and allocryptopine from *A. mexicana* that were found significantly active in a dose-dependent manner to reduce the morphine toxicity, hence suggesting that these alkaloids may be impending agents in the handling of morphine mistreatment (Capasso et al., 1997).

Lethality and safety estimate

From the seeds of *A. mexicana*, an alkaloid sanguinarine was isolated and evaluated for its hepatotoxic potential in rat model. The study revealed that a single i.p. dose (10 mg kg⁻¹) of this alkaloid, not only augmented the action of SGPT and SGOT significantly, but also created a noteworthy trouncing of P-450 (microsomal cytochrome) and the activity of benzphetamine N-methylase. In addition, the experimented rat exhibited substantial lowering in the body and liver weight, along with somewhat inflamed liver with fibrinous stuff. Further microscopic inspection of the hepatic tissue confirmed progressive cellular deterioration and cell death (Dalvil and Sanguinarine, 1985; Das et al., 2009).

Conclusion

Plants are our natural healers, and not a single plant on this earth is futile (Alam, 2019).

Nature has something in everything either explored or unexplored till date (Nooreen et al., 2018). On the basis of a number of studies, regarding this weed it is apparent that this habitually ignored, but remarkably shown plant of the arid regions remarkably has a range of benefits related to health issues beside its illegal utilization as a contaminant in mustard oil. The curative value exists in its hepato-protective nature, antimalarial, antibacterial, antiplasmodial, antiasthmatic, anti-allergic, anti-HIV properties, etc. Beside, these it has cytotoxic ability against cancerous cell lines. The plant is also used to treat, leprosy, skin diseases and to cure injuries. The plant parts also showed vaso-relaxant, larvicidal and molluscicidal action. Its phytochemical composition includes several useful alkaloids and antioxidants of pharmaceutical importance (Magaji et al., 2019). In ethnomedicinal culture, this herb has been extensively utilized to treat local infections and to mitigate the impact of high doses of morphine. Though, this weed is of common occurrence in different regions of India and abroad, its status as phytomedicinal herb yet to achieve, it is one of the least studied plants in spite of having huge medicinal value. This review recognizes the significance of *A. mexicana* somewhat equal to another member of family Papaveraceae, *Papaver somniferum* (opium poppy) to be a focus for the researcher in the area of phytomedicine/herbal formulations to add and explore this incredible plant in the floral wealth of this plant.

Acknowledgements

The author wish to acknowledge R.G.Bagdia Arts, S.B. Lakhotia Commerce and R.Bezonji Science College, Jalna.forproviding networking support through the Department of Botany.

References

Alam, A. (2019). Herbs that heal Spices: The hoard of natural remedies. *Ann. Phytomed.*, 8(2):7-18.

Anonymous (2004). In: *The Wealth of India A Dictionary of Indian Raw Materials and Industrial Products*. Council of Scientific and Industrial Research, New Delhi, 1:86-87.

Bhardwaj, D. K.; Bisht, M. S.; Jain, R. K. and Munjal, A. (1982). Phenolics from the seeds of *Argemone Mexicana*. *Phytochemistry*, 21(8):2154-2156.

Bhattacharjee, I.; Chatterjee, S.K.; Chatterjee, S. and Chandra, G. (2006). Antibacterial potentiality of *Argemonemexicana* solvent extracts against some pathogenic bacteria. *Memórias do Instituto Oswaldo Cruz*, 101(6):645-648.

Capasso, A.; Piacente, S.; Pizza, C.; De Tommasi, N.; Jativa, C. and Sorrentino, L. (1997). Isoquinoline alkaloids from *Argemonemexicana* reduce morphine withdrawal in guinea pig isolated ileum. *Planta Medica*, 63(4):326-328.

Chang, Y.C.; Chang, F.R.; Khalil, A.T.; Hsieh, P.W. and Wu, Y.C. (2003b). Cytotoxic benzophenanthridine and benzyloquinoline alkaloids from *Argemonemexicana*. *Zeitschrift für Naturforschung*, 58(7-8): 521-526.

Chang, Y.C.; Hsieh, P.W.; Chang, F.R.; Liaw, C.C.; Lee, K.H. and Wu, Y.C. (2003a). Two new properties argemexicaines A and B and the anti-HIV alkaloid 6-acetyl dihydrochelerythrine from *Argemonemexicana*. *Planta Medica*, 69:148-152.

Chitra, K.C.; Rao, S.J. and Rao, P.K. (1997). Efficacy of plant extracts for controlling cotton aphid (*Aphis gossypii*). *Indian Journal of Agricultural Sciences*, 67(3):134-135.

Dalvil, R.R. (1985). Sanguinarine: Its potential as a liver toxic alkaloid present in the seeds of *Argemonemexicana*. *Experientia*, 41(1):77-78.

Das, P.K.; Panda, P.; Pani, S.R. and Sethi, R. (2009). Hepatoprotective activity of plant *Argemonemexicana* (Linn). Against carbon tetrachloride (CCl₄) induced hepatotoxicity in rats. *International Journal of Pharmaceutical Research and Development*, 8:1-20.

Das, S. and Sukul, N.C. (1988). Nematicidal effect of the oil from the seeds of *Argemonemexicana*. *Environment and Ecology*, 6(1):194-197.

DeFilipps, R.A.; Maina, S.L. and Crepin, J. (2004). *Medicinal Plants of the Guianas (Guyana, Surinam, French Guiana)*. <http://www.mnh.si.edu/biodiversity/bdg/medicinal>

Foxcroft, L.C.; Lotter, W.D.; Runyoro, V.A. and Mattay, P.M.C. (2006). A review of the importance of invasive alien plants in the Ngorongoro Conservation Area and Serengeti National Park. *African Journal of Ecology*, 44(3):404-406.

Willcox, M.L.; Graz, B.; Falquet, J.; Sidibé, O.; Forster, M. and Diallo, D. (2007). Argemonemexicana decoction for the treatment of uncomplicated falciparum malaria. Transactions of the Royal Society of Tropical Medicine and Hygiene, 101(2): 1190-1198.